



## King's Research Portal

DOI:

[10.1016/j.neuroimage.2016.02.040](https://doi.org/10.1016/j.neuroimage.2016.02.040)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Dean III, D. C., O'Muircheartaigh, J., Dirks, H., Travers, B. G., Adluru, N., Alexander, A. L., & Deoni, S. (2016). Mapping an index of the myelin g-ratio in infants using magnetic resonance imaging. *NeuroImage*, 132(0), 225-237. <https://doi.org/10.1016/j.neuroimage.2016.02.040>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.


### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Mapping an index of the myelin g-ratio in infants using magnetic resonance imaging



Douglas C. Dean III <sup>a,\*</sup>, Jonathan O'Muircheartaigh <sup>b</sup>, Holly Dirks <sup>d</sup>, Brittany G. Travers <sup>a,c</sup>, Nagesh Adluru <sup>a</sup>, Andrew L. Alexander <sup>a,e,f</sup>, Sean C.L. Deoni <sup>d,g,h</sup>

<sup>a</sup> Waisman Center, University of Wisconsin-Madison, Madison, WI 53705, USA  
<sup>b</sup> Department of Neuroimaging, King's College London, Institute of Psychiatry, London SE5 8AF, UK  
<sup>c</sup> Department of Kinesiology, University of Wisconsin-Madison, Madison, WI 53705, USA  
<sup>d</sup> Advanced Baby Imaging Lab, Brown University School of Engineering, Providence, RI 02912, USA  
<sup>e</sup> Department of Psychiatry, University of Wisconsin-Madison, Madison, WI 53719, USA  
<sup>f</sup> Department of Medical Physics, University of Wisconsin-Madison, Madison, WI 53705, USA  
<sup>g</sup> Department of Pediatric Radiology, Children's Hospital Colorado, Aurora, CO, USA  
<sup>h</sup> Department of Radiology, University of Colorado Denver, Denver, CO, USA

## ARTICLE INFO

Article history:  
Received 7 October 2015  
Accepted 12 February 2016  
Available online 22 February 2016

## ABSTRACT

Optimal myelination of neuronal axons is essential for effective brain and cognitive function. The ratio of the axon diameter to the outer fiber diameter, known as the g-ratio, is a reliable measure to assess axonal myelination and is an important index reflecting the efficiency and maximal conduction velocity of white matter pathways. Although advanced neuroimaging techniques including multicomponent relaxometry (MCR) and diffusion tensor imaging afford insight into the microstructural characteristics of brain tissue, by themselves they do not allow direct analysis of the myelin g-ratio. Here, we show that by combining myelin content information (obtained with mcDESPOT MCR) with neurite density information (obtained through NODDI diffusion imaging) an index of the myelin g-ratio may be estimated. Using this framework, we present the first quantitative study of myelin g-ratio index changes across childhood, examining 18 typically developing children 3 months to 7.5 years of age. We report a spatio-temporal pattern of maturation that is consistent with histological and developmental MRI studies, as well as theoretical studies of the myelin g-ratio. This work represents the first ever *in vivo* visualization of the evolution of white matter g-ratio indices throughout early childhood.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

## Article info

### Recommended Articles

#### Trends and properties of human cerebral cortex: Correlations with cortical myelin content

Matthew F. Glasser, ... <sup>+3</sup> ... , David C. Van Essen  
*NeuroImage* • June 2014

📄 Preview   📄 View PDF   ⬇️ Save PDF

#### Quantitative MRI in the very preterm brain: Assessing tissue organization and myelination using magnetization transfer, diffusion tensor and T<sub>1</sub> imaging

Revital Nossin-Manor, ... <sup>+6</sup> ... , John G. Sled  
*NeuroImage* • 1 January 2013

📄 Preview   📄 View PDF   ⬇️ Save PDF

#### NODDI: Practical *in vivo* neurite orientation dispersion and density imaging of the human brain

Hui Zhang, ... <sup>+2</sup> ... , Daniel C. Alexander  
*NeuroImage* • 16 July 2012

📄 Preview   📄 View PDF   ⬇️ Save PDF

